

References

Provided by: Rebecca Andridge, Ph.D
The Ohio State University
Methods: Mind the Gap
June 17, 2022

- Andridge RR. (2011) Quantifying the impact of fixed effects modeling of clusters in multiple imputation for cluster randomized trials. *Biometrical Journal*. 2011;53(1):57–74. doi: 10.1002/bimj.201000140.
- Carpenter, J. R., Goldstein, H., & Kenward, M. G. (2011). REALCOM-IMPUTE software for multilevel multiple imputation with mixed response types. *Journal of Statistical Software*, 45(5), 1–14. doi: 10.18637/jss.v045.i05
- Drechsler, J. (2015). Multiple imputation of multilevel missing data—rigor versus simplicity. *Journal of Educational and Behavioral Statistics*, 40(1), 69–95. doi: 10.3102/1076998614563393
- Enders, C.K., Hayes, T., & Du, H. (2018). A comparison of multilevel imputation schemes for random coefficient models: Fully conditional specification and joint model imputation with random covariance matrices. *Multivariate Behavioral Research*, 53, 695-713.
- Enders, C. K., Mistler, S. A., & Keller, B. T. (2016). Multilevel multiple imputation: A review and evaluation of joint modeling and chained equations imputation. *Psychological Methods*, 21(2), 222–240. doi: 10.1037/met0000063
- Fiero, M. H., Huang, S., Oren, E., & Bell, M. L. (2016). Statistical analysis and handling of missing data in cluster randomized trials: a systematic review. *Trials*, 17, 72. <https://doi.org/10.1186/s13063-016-1201-z>

Grund, S., Lüdtke, O., & Robitzsch, A. (2018). Multiple imputation of missing data for multilevel models:

Simulations and recommendations. *Organizational Research Methods*, 21(1), 111–149.

doi: 10.1177/1094428117703686

Taljaard M, Donner A, Klar N. Imputation strategies for missing continuous outcomes in cluster

randomized trials. *Biometrical Journal*. 2008;50(3):329–45. doi: 10.1002/bimj.200710423.

Vink G, Lazendic G, van Buuren S. Partitioned predictive mean matching as a multilevel imputation

technique. *Psychol Test Assess Model*. 2015;57(4):577–94.